

ON SCHWARZSCHILD'S TOPOLOGY IN BRANE-WORLDS

E. M. MONTE*

Departamento de Fisica, Universidade
Federal da Paraiba, 58059-970, João Pessoa, Paraiba, Brazil.

and

M. D. MAIA†

Instituto de Fisica, Universidade de Brasilia, 70919-970, Brasilia,
Distrito Federal, Brazil.

February 7, 2008

Abstract

The topological structure of Schwarzschild's space-time and its maximal analytic extension are investigated in the context of brane-worlds. Using the embedding coordinates, these geometries are seen as different states of the evolution of a single brane-world. Comparing the topologies and the embeddings it is shown that this evolution must be followed by a signature change in the bulk.

1 Introduction

The emphasis in the development of higher dimensional theories has recently shifted towards the brane-world picture[1, 2]. A brane-world may be regarded as a space-time locally embedded in a higher dimensional space, the

*e-mail: edmundo@fisica.ufpb.br

†e-mail: maia@fis.unb.br

bulk, solution of higher dimensional Einstein's equations. Furthermore, the embedded geometry is assumed to exhibit quantum fluctuations with respect to the extra dimensions at the TeV scale of energies. Finally, all gauge interactions belonging to the standard model must remain confined to the four-dimensional space-time. Contrasting with other higher dimensional theories, the extra dimensions may be large and even infinite, with the possibility of being observed by TeV accelerators.

The embedding conditions relate the bulk geometry to the brane-world geometry, as it is clear from the Gauss-Codazzi-Ricci equations. Therefore, it is likely that the quantum fluctuations of the embedded geometry affect somehow the geometry of the bulk. The purpose of this paper is to show this relationship through a particular example, where we have an explicit embedding.

Since a quantum theory of embedded geometries is not yet available, it is customary to work in brane-worlds with the classical approximation, in the form of geometric perturbations. In particular, for a given embedded background geometry, we may generate a coordinate gauge independent perturbation of that background by a local shift of the embedding along the extra dimensions. The result is another geometry representing the evolution of the original brane-world[3].

In our example we take as the background a spherically symmetric brane-world (Schwarzschild). Then we perturb this geometry by extending it to the maximal analytic extension represented by Kruskal's metric. By comparing the embedding and topological differences between the background and the perturbed geometries, we conclude that the perturbation induce a change in the signature of the bulk.

Using the usual spherical coordinates (t, r, θ, ϕ) the Schwarzschild space-time is $(E, g) = ((V_4 \cup B_4), g)$, with metric g ,

$$ds^2 = (1 - 2m/r)dt^2 - (1 - 2m/r)^{-1}dr^2 - r^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (1)$$

Using its intrinsic properties, we may separate this space into two regions:[4, 5]

i) The exterior Schwarzschild space-time (V_4, g) :

$$V_4 = P_I^2 \times S^2, \quad P_I^2 = \{(t, r) \in \mathbb{R}^2 \mid r > 2m\}.$$

ii) The Schwarzschild black hole (B_4, g) :

$$B_4 = P_{II}^2 \times S^2, \quad P_{II}^2 = \{(t, r) \in \mathbb{R}^2 \mid 0 < r < 2m\}.$$

In both regions S^2 denotes the sphere of radius r .

The embeddings that are derived from the solutions of the Gauss-Codazzi-Ricci equations, without use of additional conditions are called differentiable embeddings. This is the kind of embedding that is appropriate to describe the local evolution of brane-worlds, because an otherwise rigid or an analytic embedding would be too simple or too specific to generate high frequency brane-worlds fluctuations with TeV energy, including their transient conditions.

Some relevant results are derived from differentiable embedding techniques: Collinson's theorem[6] state that no non-flat empty space-time can be embedded locally and isometrically in a five-dimensional space of nonzero constant curvature. A theorem due to Szekeres tells that no non-flat vacuum metric can be embedded in five dimensions[7]. Therefore, the Schwarzschild's background cannot be differentiably embedded in a five dimensional flat or constant curvature bulk, as in the Randall-Sundrum model. For simplicity, we will consider the case of six-dimensional flat bulks $M_{r,s}$ with signatures $(r, s) = (4, 2)$ and $(r, s) = (5, 1)$, where the Schwarzschild space-time is known to be embeddable.

2 The Embedding Structure

The two known local immersions of the Schwarzschild space-time into six dimensional pseudo-Euclidean bulks are: (using $2m = 1$)

The Kasner embedding [8]

$$Y : (E, g) \rightarrow M_{4,2} \left\{ \begin{array}{l} Y_1 = (1 - 1/r)^{1/2} \cos t \\ Y_2 = (1 - 1/r)^{1/2} \sin t \\ Y_3 = f(r), \quad (df/dr)^2 = \frac{1+4r^3}{4r^3(r-1)} \\ Y_4 = r \sin \theta \sin \phi \\ Y_5 = r \sin \theta \cos \phi \\ Y_6 = r \cos \theta \end{array} \right\}$$

and the Fronsdal embedding [9]

$$Y' : (E, g) \rightarrow M_{5,1} \left\{ \begin{array}{l} Y'_1 = 2(1 - 1/r)^{1/2} \sinh(t/2) \\ Y'_2 = 2(1 - 1/r)^{1/2} \cosh(t/2) \\ Y'_3 = g(r), \quad (dg/dr)^2 = \frac{(r^2+r+1)}{r^3} \\ Y'_4 = r \sin \theta \sin \phi \\ Y'_5 = r \sin \theta \cos \phi \\ Y'_6 = r \cos \theta \end{array} \right\}.$$

Notice that Y'_3 is defined for $r > 0$, while Y_3 is defined only for $r > 1$, suggesting the extension of $(E, g) = ((V_4 \cup B_4), g)$.

Now we can show that the Fronsdal embedding indeed corresponds to an extension of the Schwarzschild metric. For this purpose, define the new coordinates u and v by: For $r > 2m$,

$$v = \frac{1}{4m} \left(\frac{r}{2m} \right)^{1/2} \exp\left(\frac{r}{4m}\right) Y'_1 \quad \text{and} \quad u = \frac{1}{4m} \left(\frac{r}{2m} \right)^{1/2} \exp\left(\frac{r}{4m}\right) Y'_2. \quad (2)$$

For $0 < r < 2m$,

$$v = \frac{i}{4m} \left(\frac{r}{2m} \right)^{1/2} \exp\left(\frac{r}{4m}\right) Y'_1 \quad \text{and} \quad u = \frac{i}{4m} \left(\frac{r}{2m} \right)^{1/2} \exp\left(\frac{r}{4m}\right) Y'_2, \quad (3)$$

where

$$u^2 - v^2 = \left(\frac{r}{2m} - 1 \right) \exp\left(\frac{r}{2m}\right) \iff Y'^2_2 - Y'^2_1 = 16m^2 \left(1 - \frac{2m}{r} \right). \quad (4)$$

Now $r = r(Y'_1, Y'_2)$ is implicitly defined by equation (4), while $t = t(Y'_1, Y'_2)$ is implicitly defined by

$$Y'_1/Y'_2 = \tanh\left(\frac{t}{4m}\right). \quad (5)$$

Finally, the metric g' in the new coordinates is

$$ds^2 = (32m^3/r) \exp(-r/2m)(dv^2 - du^2) - r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad (6)$$

which is the same metric encountered by Kruskal[10] representing the maximal analytic extension of the Schwarzschild metric without a singularity at $r = 2m$. Using the new coordinates (u, v, θ, ϕ) , the Schwarzschild space-time (E, g) is extended to space-time $(E' = Q^2 \times S^2, g')$, where Q^2 is the Kruskal plane defined by (4).

3 The Topological Structure

In the following we show that the topology of (E, g) is $(\{\mathbb{R}^2 - \{(t, r) \in \mathbb{R}^2 \mid r = 2m\}\} \times S^2)$ and that the topology of (E', g') is $\mathbb{R}^2 \times S^2$, when the space (E, g) is embedded in different bulks. Indeed, by construction: $E = [P_I^2 \cup P_{II}^2] \times S^2$ and $E' = Q^2 \times S^2$. The topology of E is the Cartesian product topology of $[P_I^2 \cup P_{II}^2]$ by S^2 , while that the topology of E' is the Cartesian product topology of Q^2 by S^2 . While the topology $S^2 \subset \mathbb{R}^3$ is the usual topology[11] induced by the topological space $(\tau_3, \mathbb{R}^3)$, the topologies of $[P_I^2 \cup P_{II}^2] \subset \mathbb{R}^2$ and of $Q^2 \subset \mathbb{R}^2$, respectively, τ_p and τ_q , are induced from $(\tau_2, \mathbb{R}^2)$.

Since Q^2 is an extension of $[P_I^2 \cup P_{II}^2]$, we may define the embedding, $\psi : [P_I^2 \cup P_{II}^2] \rightarrow Q^2$. Therefore, defining the open set $A = \{(t, r) \in \mathbb{R}^2 \mid t^2 + (r - 2m)^2 < m^2 \text{ and } r > 0\}$, we have $A \cap [P_I^2 \cup P_{II}^2] = A - \{(t, r) \in \mathbb{R}^2 \mid r = 2m\}$. This is an open set of the topological space $[P_I^2 \cup P_{II}^2]$, composed of two connected components which form a topological basis for the semi-plane $t - r, r > 0$.

On the other hand, we have that $\psi(A \cap [P_I^2 \cup P_{II}^2])$ is given for another open set composed by four connected components. Using (4) it follows that the lines L_1 and L_2 defined corresponding to $r = 2m$ are on Q^2 , we have that $\{\psi(A \cap [P_I^2 \cup P_{II}^2]) \cup L_1 \cup L_2\} \cap D = B$, where D is an open disk on $\mathbb{R}^2$ with center in the origin of Q^2 . Therefore, B is a plane disk in the new coordinates $r = r(Y'_1, Y'_2)$ and $t = t(Y'_1, Y'_2)$. Consequently, the topology of Q^2 is given by open sets of $\mathbb{R}^2$.

Our conclusion is that the topology of (E, g) , $(\mathbb{R}^2 - \{(t, r) \in \mathbb{R}^2 \mid r = 2m\}) \times S^2$, is different from the topology $\mathbb{R}^2 \times S^2$ of the extension (E', g') . Notice that (E, g) is composed by two components and consequently it is

disconnected, while (E', g') is connected as it follows from the Fronsdaal embedding.

While the embedding of the Schwarzschild space-time given by Kasner remains differentiable only for $r > 2m$, or in another connected component $0 < r < 2m$, the Fronsdaal embedding remains differentiable all the way to $r = 0$ (for $r > 0$ in a only connected component). If the latter is regarded as an evolution of the former, then this is only possible with a change of signature and topology of bulk. These changes should be observable at the predicted laboratory production of black-holes.

References

- [1] Nima Arkani-Hamed *et al.*, *Phys. Lett.* **B429**, 263 (1998).
- [2] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83**, 3370 (1999).
- [3] M. D. Maia and E. M. Monte, *Phys. Lett.* **A297**, 9 (2002).
- [4] B. O'Neill, *Semi-Riemannian Geometry*, Academic Press, New York, (1983).
- [5] M. D. Maia and E. M. Monte, *Contemp. Math.*, **13**, 229 (1997).
- [6] C. D. Collinson, *J. Phys.* **A4**, 206 (1971).
- [7] P. Szekeres, *Nuovo Cimento*, **43**, 1062 (1966).
- [8] E. Kasner, *Am. J. Math.* **43**, 130 (1965).
- [9] C. Fronsdaal, *Phys. Rev.* **116**, 778 (1959).
- [10] M. Kruskal, *Phys. Rev.* **119**, 1743 (1960).
- [11] J. Dugundji, *Topology*, Ally and Bacon Series in Advanced Mathematics, Boston, (1966).
- [12] V. A. Rubakov, hep-ph/0104152.
- [13] K. Maeda *et al.*, *Phys. Rev.* **D62**, 24012, (2000).